

“New” marine processes and features observed at <1 - 10 m resolution

**Quinten Vanhellemont
RBINS/DO Nature**

Presented at IOCS 16 June 2015

<http://odnature.naturalsciences.be/remsem/>



Landsat-8 « Gallery »

NW Australia – 2013-05-26
Landsat-8 / RBINS processing

15°4'S
15°8'S
15°12'S
15°16'S

124°28'E


124°32'E

124°36'E

124°40'E

124°44'E

10 km




NW Australia – 2013-06-11
Landsat-8 / RBINS processing

15°4'S
15°8'S
15°12'S
15°16'S

124°28'E 124°32'E 124°36'E 124°40'E 124°44'E

10 km




Alaska – 2014-09-06

Landsat-8 / RBINS processing



Tunisia/Lake Ichel and Lake Bizerte – 2013-08-28

Landsat-8 / RBINS processing



Tunisia/Lake Ichel and Lake Bizerte – 2014-03-24
Landsat-8 / RBINS processing



Land

GB

51°20'N 51°24'N 51°28'N 51°32'N

51°23'N 51°24'N 51°25'N 51°26'N 51°27'N

2°56'E

3°22'E

3°24'E


3°26'E

3°28'E

2 km

white wake (foam)

brown wake (sediment)





Gunfleet Sands


London Array

**Thames Estuary – 2014-08-21
Landsat-8 / RBINS processing**



Venezuela/Lake Maracaibo– 2014-02-27
Landsat-8 / RBINS processing

Gulf of Mexico Oil Rigs - 2015-02-23
Landsat-8 / RBINS processing



A satellite image showing several oil rigs in the Gulf of Mexico. The water is dark blue. Several bright lights from the rigs are visible, along with plumes of white smoke and blue oil slicks on the water's surface.

gas flares

smoke

oil slicks?

51°19'N 51°20'N 51°21'N 51°22'N 51°23'N 51°24'N

Belgium/AOC Zeebrugge – 2013-08-04
Landsat-8 / RBINS processing

Aeronet-OC MOW1



3°2'E

3°4'E

3°6'E

3°8'E

3°10'E

3°12'E

51°19'N 51°20'N 51°21'N 51°22'N 51°23'N 51°24'N

Belgium/AOC Zeebrugge – 2013-10-07
Landsat-8 / RBINS processing

Aeronet-OC MOW1



3°2'E

3°4'E

3°6'E

3°8'E

3°10'E

3°12'E



Australia/AOC Lucinda – 2013-06-25
Landsat-8 / RBINS processing



Aeronet-OC
Lucinda

Sweden/AOC Gustav Dalen – 2014-07-21


Landsat-8 / RBINS processing

Aeronet-OC Gustav Dalen





Landsat-8 « Applications »

Belgian/Dutch coastal zone and Western Scheldt Estuary





Landsat-8/OLI Rayleigh corrected RGB 2014-09-08

Turbidity (FNU)




Landsat-8/OLI Turbidity (V2015+D2015) 2014-09-08

Turbidity (FNU)




Landsat-8/OLI Turbidity (V2015+D2015) 2014-09-08

London Array offshore wind farm





Vanhellemont, Q., Ruddick, K., 2014a. **Turbid wakes associated with offshore wind turbines observed with Landsat 8**. *Remote Sens. Environ.* 145, 105–115.

OLI-derived suspended particulate matter (SPM)




Vanhellemond, Q., Ruddick, K., 2014a. **Turbid wakes associated with offshore wind turbines observed with Landsat 8.** *Remote Sens. Environ.* 145, 105–115.

Gunfleet Sands wind farm (UK)



Landsat-8/OLI 2013-10-30

Rayleigh corrected RGB



Vanhellemont, Q., Ruddick, K., 2014b. **Landsat-8 as a Precursor to Sentinel-2: Observations of Human Impacts in Coastal Waters.**, in: ESA Special Publication SP-726.

Simple detection algorithm for high concentrations of black sediments



the NIR marine reflectance is greater than a threshold for turbid waters:

$$\rho_w^5 > 0.01$$

the maximum marine reflectance in three visible bands is low for this turbidity:

$$\max(\rho_w^2, \rho_w^3, \rho_w^4) < 0.07$$

L8/OLI 2014-03-16



Landsat-8/OLI 2013-10-30



Vanhellemont, Q., Ruddick, K., (2015a). Advantages of high quality SWIR bands for ocean colour processing: examples from Landsat-8. <http://dx.doi.org/10.1016/j.rse.2015.02.007>


Landsat-8/OLI 2013-10-30

$\rho_w^5 > 0.01$

$\max(\rho_w^2, \rho_w^3, \rho_w^4) < 0.07$


$\rho_w^5 > 0.001 \approx \text{ambient}$

$\max(\rho_w^2, \rho_w^3, \rho_w^4) < 0.07$



Higher resolution?

Pan-sharpening for marine applications (details on poster)




(a) 30 m

ship

1 km

(b) 15 m sharpened



surface gravity waves

sharper features

Pléiades very high resolution (2.8 m MS, 0.7 PAN)





A satellite image showing a coastal area. The left side is dominated by green land with some brownish patches, likely indicating agricultural fields or coastal erosion. A large, light brown body of water occupies the right side, representing a harbor entrance. A small, dark, elongated object, possibly a boat or a bridge, is visible in the lower right corner of the water area.

Zeebrugge – harbour entrance (PANs)
Pléiades 2014-09-08 – RBINS processing

www.deme-group.com



Zeebrugge – dredger (PANs)
Pléiades 2014-09-08 – RBINS processing

Zeebrugge – patchy SPM (PANs)
Pléiades 2014-09-08 – RBINS processing



Patchy surface suspended sediment concentration
RV Belgica campaign April 2015

building
shadow →




Blankenberge yacht harbour (PANs)


Pléiades 2014-09-08 – RBINS processing

Zeebrugge – MOW1 tidal wake (PANs)
Pléiades 2014-09-08 – RBINS processing








Pléiades Rayleigh corrected RGB 2014-09-08




Pléiades Turbidity (ACOLITE a/c + D2015) 2014-09-08





Pléiades 2014-09-08 11:10 UTC




C-Power Offshore Transformer Station (PANs)
Pléiades 2015-04-14 / RBINS processing



Aeronet-OC





Platform impact on moderate resolution pixel (see poster)

Landsat-8/OLI data

